



# Abrikossoff tumour mimicking cecal polyp – case report and literature review of granular cell tumours the of gastrointestinal tract

Anna Rycyk-Bojarzyńska<sup>1,B,D</sup>✉, Kinga Knop-Chodyła<sup>2,C-D</sup>, Agnieszka Szczerbińska<sup>3,A,D</sup>,  
Beata Kasztelan-Szczerbińska<sup>1,B,E</sup>, Halina Cichoż-Lach<sup>1,B,F</sup>, Dorota Lewkowicz<sup>4,B-C</sup>,  
Justyna Szumiło<sup>4,B-C</sup>

<sup>1</sup> Department of Gastroenterology with Endoscopy Unit, Medical University, Lublin, Poland

<sup>2</sup> Department of Gastroenterology with Endoscopy Unit, University Clinic Hospital Number 4, Lublin, Poland

<sup>3</sup> Student Campus, Medical University, Warsaw, Poland

<sup>4</sup> Department of Pathomorphology, Medical University, Lublin, Poland

A – Research concept and design, B – Collection and/or assembly of data, C – Data analysis and interpretation, D – Writing the article, E – Critical revision of the article, F – Final approval of the article

Rycyk-Bojarzyńska A, Knop-Chodyła K, Szczerbińska A, Kasztelan-Szczerbińska B, Cichoż-Lach H, Lewkowicz D, Szumiło J. Abrikossoff tumour mimicking cecal polyp – a case report and literature review of granular cell tumours the of gastrointestinal tract. *Ann Agric Environ Med*. doi: 10.26444/aaem/192795

## Abstract

Granular cell tumour (GCT) or Abrikossoff tumour, is a rare submucosal neoplasm originating from Schwann cells, and accounts for 0.03% of all submucosal tumours. Only 8% of GCTs occur in the gastrointestinal tract, mostly in the esophagus. Diagnosis of granular cell tumour is exceptionally rare. The case is presented of a 72-year-old female patient who underwent polypectomy of a colon polyp which, in the histopathology report, appeared to be a granular cell tumour. The possibility of granular cell tumour diagnosis which is extremely rare, is discussed. However, 2% of these tumours are malignant. It is very important to establish the diagnosis as early as possible. The patient was admitted to the Gastroenterology Outpatient Clinic due to a hepatic vascular lesion. She did not present any symptoms of granular cell tumour. A complete resection of the granular cell tumour fortunately confirmed it to be benign.

## Key words

cancer, tumour, colon, polyp, granular cell tumour (GCT), Abrikossoff tumour, Abrikosov's tumour

## INTRODUCTION

Granular cell tumours (GCTs) are rare soft tissue neoplasms that can affect virtually any organ and any age group in patients [1]. They are most commonly localized in the oral cavity, skin and subcutaneous tissue. Their occurrence in the gastrointestinal (GI) tract comprises only 8% of all GCTs [2]. They are diagnosed in patients most often between the fourth and sixth decades of life, with a female-to-male ratio of 2:1 [3]. Most lesions are benign, but 1–2% are diagnosed as malignant, which is associated with a worse prognosis [4, 5]. The case is presented of a 72-year-old female patient who underwent colonoscopy due to the high risk of colon cancer in her age group. A polypectomy of a colon polyp was performed and the histopathology report confirmed the presence of GCT.


## CASE REPORT

The female patient presented to the Gastro-enterology Outpatient Clinic due to a hepatic vascular lesion found incidentally on abdominal computed tomography (CT) scan. She denied any complaints of abdominal pain, although having had a history of left mastectomy for breast cancer

with no follow-up treatment. In control mammography, no changes were detected for further observation. The patient additionally suffers from chronic ischemic heart disease, hypertension, paroxysmal atrial fibrillation, diabetes mellitus, and depression. Due to years of nicotine use and diagnosed chronic obstructive pulmonary disease (COPD), she remains under the control of a pulmonology clinic. Recently, there has been an increase in persistent cough. The pulmonologist ordered a follow-up chest X-ray, which showed lesions of an unclear nature and dilatation of the hilar shadows. Due to the indeterminate nature of the lesions, a review CT scan of the chest and abdominal cavity was performed. Examination showed emphysema, atelectatic bilateral consolidation of the lung parenchyma, and multiple lesions of a calcified nodule nature, ranging from 2mm – 8mm in diameter. Within the abdominal cavity, foci of contrast enhancement with features of vascular malformation in the liver were detected. The hepatic lesion prompted the extension of further gastro-enterological diagnosis to endoscopic studies. Esophagogastroduodenoscopy revealed features of inflammation in the antrum, angle and body of the stomach which were confirmed in histopathological examination. In Warthin-Starry staining, in the mucus on the surface of the stomach, numerous bacteria corresponding to *Helicobacter pylori* infection were found. The patient was recommended to undergo *H. Pylori* eradication. During colonoscopy, a cecal polyp was removed by endoscopic submucosal dissection.

Histopathological examination of the removed polyp revealed a granular cell tumour – identified as an Abrikossoff

✉ Address for correspondence: Anna Rycyk-Bojarzyńska, Department of Gastroenterology with Endoscopy Unit, Medical University, Lublin, Poland  
E-mail: aniarycyk@op.pl



**Figure 1.** Microscopic appearance of caecal granular cell tumour. Sheets of large polygonal cells with abundant granular cytoplasm and small dense nuclei with moderate atypia (A, B). Strongly positive cytoplasmic immunostaining for S-100 in neoplastic cells (C). (A, B – HE, C – IHC; objective magnification A, C – 10x, B – 20x)

tumour; however, the margins of the resected polyp were free of tumour. On immunohistochemical analysis, there was positive staining for S-100 protein expression, and negative for desmin.

The patient remains under the control of the Gastroenterology Outpatient Clinic. The patient was planned to follow-up colonoscopy in one year. Subsequent decisions regarding surveillance will be made based on colonoscopy results. At present, the patient presents no symptoms in the gastrointestinal tract.

## DISCUSSION AND CONCLUSIONS

**Occurrence.** Granular cell tumour (GCT) is a lesion of controversial origin. It was first classified as a myogenic tumour in 1926 by the Russian pathologist Abrikossoff [5]. However, it was only with the development of immunohistochemistry and electron microscopy techniques that it was properly determined to originate from nerve tissue-Schwann cells [6]. Currently, the etiology of granular cell tumours and their genetic basis is still poorly understood. However, studies show that the tumour often co-occurs with genetic syndromes, such as LEOPARD syndrome or Noonan syndrome [7]. Whole genome sequencing capabilities and a 2018 study by Pareja et al. also identified recurrent somatic

mutations inactivating ATP6AP1 and ATP6AP2 in 72% of granular cell tumours [8]. Due to its as yet incompletely understood origin and rare occurrence, epidemiological data will begin to be limited. Knowledge of granular cell tumour originates mainly from clinical case reports [9]. GCTs can appear at any age [10], and the lesions can affect almost any organ. Their frequency is shown in Table 1 [2–3,5,7,11–13]. However, the following data remain approximate, owing to the different values presented in literature reviews.

**Table 1.** The most common locations of granular cell tumours along with the incidence

Location	Prevalence (%)
Oral cavity [2, 5, 7, 11–12]	40%
Skin and subcutaneous tissue [2, 5, 7, 11–13]	30%
Breast [2, 5, 7, 11]	15%
Respiratory tract [2, 5, 7, 11–12]	10–15%

Only 8% of GCTs occur in the gastrointestinal tract, with the most common cases in the esophagus, large intestine and stomach [2]. Only 150 cases of GCTs in the large intestine have been reported in the literature [14]. The most common locations of GCTs of the gastrointestinal tract are shown in Table 2.

**Table 2.** Frequency of granular cell tumours in different locations of the gastrointestinal tract [1,5,10,14–19]

Location	Prevalence (%)
Oesophagus	65–68%
Colorectal	20–21%
Stomach	4–9%
Small intestine	~1%
Anal	~1%

**Symptoms.** The symptoms described below relate to locations in the gastrointestinal tract only. Most often, granular cell tumours do not produce clinical symptoms and their appearance depends on the size of the tumour. They are often detected incidentally in endoscopy procedures performed for other reasons. The situation was similar in the patient in the described case report. If present, symptoms are non-specific and depend on the location of the lesion. In the oesophagus, GCT can manifest as gastroesophageal reflux or dysphagia. In the small and large intestine, it can show bleeding or abdominal pain, and in the large intestine, it can additionally cause a change in the nature of bowel movements [1,16].

**Diagnosis.** The main diagnostic method for GCT is gastrointestinal endoscopy. Mostly, colonoscopy reveals GCT as a yellowish-white colour sitting polyps or nodules. This is commonly the case due to the fact that GCTs have no specific features that distinguish them from other intestinal polyps [14]. Only 7–25% of GCTs in this area occur in multiple forms [16]. Colorectal GCTs are mostly located in the ascending colon and cecum [17].

**Histological findings.** Histopathological and immunohistochemical examination remains the standard for the diagnosis of granular cell tumours [14]. Morphologic features include nests of polygonal, spindle-shaped cells with abundant eosinophilic, fine-grained cytoplasm [1]. A typical histologic feature of GCTs is the pustule-ovoid bodies of Milian (POBoM), which are detected under microscope as a pale halo surrounding the eosinophilic bodies [20]. They usually show positive staining for S-100, CD68, neuron-specific enolase, CD57, inhibin, TFE3, SOX10, CD56 and vimentin [15,21]. However, there is sometimes a variant that is negative for S-100, but remains positive for CD68, CD10, and sometimes neuron-specific enolase [22].

Although most GCTs are benign lesions, on the basis of histologic features, in 1998, Fanburg-Smith et al. proposed distinguishing features based on which lesions can be divided into benign, atypical and malignant. These include histological features: necrosis, spindling, vesicular nuclei with large nucleoli, and increased mitotic count [23]. Interestingly, none of them were detected in the presented case.

Some lesions are routinely removed by polypectomy followed by histopathological examination, which occurred in the presented patient.

**Treatment.** Currently, there are no established guidelines for the treatment of granular cell tumours [1, 18–19]. However, endoscopic resection of these tumours with negative margins is recommended by most authors because GCTs can have a possible aggressive biological potential [12]. The mainstay is excision of the lesion by endoscopic mucosal

resection (EMR), endoscopic submucosal dissection (ESD) or polypectomy [16, 24]. For lesions smaller than 2cm, EMR or ESD remains preferred, for lesions 2–3cm ESD and for lesions larger than 3cm, surgical excision or consideration of ESD is advised [1, 24]. Currently, radiotherapy and chemotherapy are not recommended for patients with granular cell tumour, but studies are underway on the effect of Pazopanib for metastatic lesions [25]. The prognosis of patients with complete removal of a lesion histopathologically determined to be benign is very good. In contrast, for malignant GCTs, the probability of recurrence and 5-year survival rate is 60–75% [13].

## CONCLUSIONS

Although GCT is relatively rare in the colon, it should be considered in the differential diagnosis of submucosal lesions. It also remains important to recognize potentially malignant features of the neoplasm to implement appropriate monitoring and diagnostic vigilance especially in this group of patients.

## REFERENCES

- Barakat M, Kar AA, Poursahid S, et al. Gastrointestinal and biliary granular cell tumour: diagnosis and management. *Ann Gastroenterol*. 2018 Jul-Aug;31(4):439–447. doi: 10.20524/aog.2018.0275.
- Ramai D, Lai J, Changela K, Anand S. Colonic Granular Cell Tumour: An Endoscopic and Histopathologic Review with Case Illustration. *Cureus*. 2018 Jan 2;10(1):e2015. doi: 10.7759/cureus.2015.
- Rajagopal MD, Gochhait D, Shanmugan D, Barwad AW. Granular Cell Tumour of Cecum: A Common Tumour in a Rare Site with Diagnostic Challenge. *Rare Tumours*. 2017 Aug 29;9(2):6420. doi: 10.4081/rt.2017.6420.
- Shah K, Fulbright J, Katz A, Zhang L, Jarka D, Lypka M. Malignant Granular Cell Tumour of the Lower Extremity in an Adolescent Male. *Journal of Pediatric Hematology/Oncology*. 2021;43(4):e529. doi:10.1097/MPH.0000000000001886.
- Salaouatchi MT, De Breucker S, Rouvière H, et al. A Rare Case of a Metastatic Malignant Abrikossoff Tumour. *Case Rep Oncol*. 2021;14(3):1868–1875. doi:10.1159/000520385.
- Rekhi B, Jambhekar NA. Morphologic spectrum, immunohistochemical analysis, and clinical features of a series of granular cell tumours of soft tissues: a study from a tertiary referral cancer center. *Ann Diagn Pathol*. 2010 Jun;14(3):162–7. doi: 10.1016/j.anndiagpath.2010.01.005.
- Neelon D, Lannan F, Childs J. Granular Cell Tumour. 2023 Jul 3. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan–.
- Pareja F, Brandes AH, Basili T, et al. Loss-of-function mutations in ATP6AP1 and ATP6AP2 in granular cell tumours. *Nat Commun*. 2018 Aug 30;9(1):3533. doi: 10.1038/s41467-018-05886-y.
- Cantella R, Evola G, Di Stefano C, Trusso Zirna E, Iudica M, Piazza L. Guz ziarnistej komórki jelita ślepego: Opis przypadku mini inwazyjnej resekcji chirurgicznej i przegląd literatury. *Int J Surg Case Rep*. 2021 Oct;87:106397. doi: 10.1016/j.ijscr.2021.106397. Epub 2021 13 września.
- Gao S, Xing B, Lichao S, et al. Malignant granular cell tumour in the thoracic wall: A case report. *Front Oncol*. 2022;12:895924. doi:10.3389/fonc.2022.895924.
- Sunkara T, Nagaraj SV, Gaduputi V. Granular Cell Tumour of Rectum: A Very Rare Entity. *Case Rep Gastrointest Med*. 2017;2017:3795482. doi: 10.1155/2017/3795482. Epub 2017 Jan 31.
- Jobrack AD, Goel S, Cotlar AM. Granular Cell Tumour: Report of 13 Cases in a Veterans Administration Hospital. *Military Medicine*. 2018;183(9–10):e589–e593. doi:10.1093/milmed/usx237.
- Moten AS, Zhao H, Wu H, Farma JM. Malignant granular cell tumour: Clinical features and long-term survival. *J Surg Oncol*. 2018 Nov;118(6):891–897. doi: 10.1002/jso.25227. Epub 2018 Sep 9.
- Dhruv S, Atodaria KP, Gurala D, El Imad T, Abergel J. Granular Cell Tumour of the Ascending Colon. *Case Rep Gastroenterol*. 2023 Feb 17;17(1):104–108. doi: 10.1159/000529170.

15. An S, Jang J, Min K, et al. Granular cell tumour of the gastrointestinal tract: histologic and immunohistochemical analysis of 98 cases. *Hum Pathol.* 2015 Jun;46(6):813–9. doi: 10.1016/j.humpath.2015.02.005.
16. Guzman Rojas P, Robalino Gonzaga ES, Zayat V, Parikh J. Benign Granular Cell Tumour of the Cecum. *Cureus.* 2019 Feb 14;11(2):e4074. doi: 10.7759/cureus.4074.
17. Kawashima K, Hikichi T, Onizawa M, et al. Colonic Endoscopic Submucosal Dissection for a Granular Cell Tumour with Insufficient Endoscopic Manipulation in the Hepatic Flexure. *Case Rep Gastroenterol.* 2022;16(1):216–222. doi:10.1159/000523963.
18. Kim IK, Ju YT, Kim HG, et al. Experience of surgical treatment in a granular cell tumour in the descending colon: a case report. *Ann Coloproctol.* 2023;39(3):275–279. doi:10.3393/ac.2020.00836.0119.
19. Nitta T, Ohta M, Kataoka J, et al. Granular cell tumour coexisting with adenocarcinoma in the stomach: Report of a rare case. *Ann Med Surg (Lond).* 2021;65:102271. doi:10.1016/j.amsu.2021.102271.
20. Kent T, Saulino D. Incidental multifocal granular cell tumour in the setting of chronic gastritis discovered following gastric sleeve operation: a case report and brief review of the literature. *J Surg Case Rep.* 2024;2024(1):rjad700. doi:10.1093/jscr/rjad700.
21. Schoolmeester JK, Lastra RR. Granular cell tumours overexpress TFE3 without corollary gene rearrangement. *Hum Pathol.* 2015 Aug;46(8):1242–3. doi: 10.1016/j.humpath.2015.04.004.
22. Fernandez-Flores A, Cassarino DS, Riveiro-Falkenbach E, et al. Cutaneous dermal non-neural granular cell tumour is a granular cell dermal root sheath fibroma. *J Cutan Pathol.* 2017 Jun;44(6):582–587. doi: 10.1111/cup.12926.
23. Shrestha B, Khalid M, Gayam V, et al. Metachronous Granular Cell Tumour of the Descending Colon. *Gastroenterology Res.* 2018;11(4):317–320. doi:10.14740/gr1045w.
24. Take I, Shi Q, Qi ZP, et al. Endoscopic resection of colorectal granular cell tumours. *World J Gastroenterol.* 2015 Dec 28;21(48):13542–7. doi: 10.3748/wjg.v21.i48.13542.
25. Wei L, Liu S, Conroy J, et al. Whole-genome sequencing of a malignant granular cell tumour with metabolic response to pazopanib. *Cold Spring Harb Mol Case Stud.* 2015 Oct;1(1):a000380. doi: 10.1101/mcs.a000380.